

FOSSIL ENERGY, CO₂, CLIMATE CHANGE, AND THE AEROSOL PROBLEM

Stephen E. Schwartz

For presentation at the
2011 IYC O₃ Symposium on Stratospheric Ozone and Climate Change,
Washington, DC
Nov. 7-10, 2011

**Environmental Sciences Department/Atmospheric Sciences Division
Brookhaven National Laboratory**

**U.S. Department of Energy
Office of Science**

ABSTRACT

Climate change due to increasing atmospheric carbon dioxide differs from stratospheric ozone depletion, regional air pollution, and acid deposition, because emission of CO₂ is intrinsically coupled to production of the energy on which our industrial society relies. Unlike chlorofluorocarbons there are no readily available substitutes, and unlike sulfur or nitrogen oxides and hydrocarbons, CO₂ can be removed from the waste stream only with difficulty and great expense (energy penalty). Consequently, until alternative energy sources are developed or a practical means of sequestering CO₂ is developed, we are faced with continued CO₂ emissions. Determining how much more CO₂ can be emitted into the atmosphere consistent with a given allowable increase in global temperature above preindustrial, e.g., 2°C, requires knowledge of Earth's climate sensitivity, the increase in global temperature per increase in radiative forcing, which is uncertain by about a factor of 2. The observed increase in temperature over the twentieth century is well less than that which would be expected from radiative forcing by incremental greenhouse gases alone together with the current best estimates of climate sensitivity and planetary energy imbalance. Possible explanations are a lower climate sensitivity and/or partial offset of the greenhouse gas forcing due to cooling forcing by tropospheric aerosols; as aerosols, unlike CO₂, are short-lived in the atmosphere, reduction of emission of aerosols and precursor gases would reveal the committed warming that has been masked by the aerosol cooling forcing. Allowable future CO₂ emissions so as not to commit the planet to a temperature increase of 2°C above preindustrial are at most a few decades at present emission rates; the possibility that Earth is already committed to such a temperature increase cannot be ruled out. These considerations conspire to make climate change a challenging scientific problem and an enormously difficult societal issue.

NOTICE: This manuscript has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-98CH10886 with the U.S. Department of Energy. The publisher by accepting the manuscript for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes.